

Quzyttir[®]

Cetirizine HCl Injection
10 mg/mL



Interested in another treatment option for your short-term hives?

This is your Q
to ask about
next-generation
IV antihistamine
QUZYTIR

Actor portrayal.

What is QUZYTIR?

QUZYTIR is an FDA-approved prescription medicine for the treatment of short-term hives (known as acute urticaria) in adults and children 6 months of age and older.

QUZYTIR is not recommended in children younger than 6 years old with kidney or liver problems.

SELECT SAFETY INFORMATION

- You should not receive QUZYTIR if you are allergic to any of its ingredients or if you are allergic to other medicines that contain levocetirizine or hydroxyzine.

Please see additional Important Safety Information on back.

Please see accompanying or click for [Full Prescribing Information](#).

Antihistamines are important in the treatment of short-term hives, a common symptom of allergic reactions

What are antihistamines?

An allergist or doctor will likely recommend an antihistamine for your hives. This treatment works by blocking the effect of histamine, a chemical released by the immune system that can cause hives and other allergy symptoms.

Are there different types of antihistamines?

Yes, there are two generations of antihistamines. Both are effective in the treatment of hives, but there are differences.

- **First-generation** antihistamines, like **diphenhydramine**, are known to have more side effects, such as drowsiness, and **more interactions with other medications**
- **Second-generation** antihistamines were made to **improve on side effects like drowsiness** and have **no major interactions with other medications**

Tell your healthcare professional about all medicines you take, including prescription or over-the-counter medicines, vitamins, or herbal supplements.

Discover next-generation QUZYTTR

The first and only second-generation intravenous (IV) antihistamine

QUZYTTR is an FDA-approved prescription medicine for the treatment of short-term hives (known as acute urticaria) in adults and children 6 months of age and older. It is not recommended in children younger than 6 years old with kidney or liver problems.

How is QUZYTTR administered?

QUZYTTR is a **single-use vial** that must be administered intravenously by a medical professional. It is preservative-free and **does not require refrigeration**.

The recommended dosage regimen is **once every 24 hours** as needed for symptom relief. The recommended dosage depends on your age and doctor's judgment.

Where is QUZYTTR administered?

You may receive QUZYTTR in your **doctor's office, at your infusion center, in urgent care clinics, or while at the hospital**. If needed, some people may receive at-home treatment by a medical professional.



Not actual size.

SELECT SAFETY INFORMATION

- QUZYTTR may cause sleepiness. Be careful when driving a car or operating potentially dangerous machinery. Avoid alcohol or taking certain other medicines with QUZYTTR, as this may raise your risk of having problems with alertness or thinking. Tell the healthcare provider treating you about all the medicines you are taking, including prescription or over-the-counter medicines, vitamins, or herbal supplements.

Please see additional Important Safety Information on back.

Please see accompanying or click for [Full Prescribing Information](#).

Learn how QUZYTIR (cetirizine hydrochloride injection) can help

In a clinical study, patients were given either QUZYTIR (n=127) or IV diphenhydramine (n=135) for their hives.

Patients **rated their level of itch from hives** before treatment and 2 hours after treatment with either QUZYTIR or IV diphenhydramine:



QUZYTIR worked like IV diphenhydramine at reducing hives from allergic reactions.

Patients were also followed to see **if their hives returned** and if they needed additional care:

2X

fewer return visits

Less than half of **QUZYTIR** patients had to return after discharge

6%

of **QUZYTIR** patients had to return

14%

of **IV diphenhydramine** patients had to return

SELECT SAFETY INFORMATION

- QUZYTIR was not studied in pregnant or nursing women. Tell your healthcare provider if you are pregnant, plan to become pregnant, or are breastfeeding.

In the same clinical study, **patients rated their level of drowsiness** with either QUZYTTIR or IV diphenhydramine.



Less drowsiness

QUZYTTIR patients were not as tired during and after treatment

How does QUZYTTIR work differently?



Starts fast

Starts working in less than 4 minutes



Lasts long

Lasts for 24 hours



Actor portrayal.

SELECT SAFETY INFORMATION

- The most common side effects (which occurred in <1% of patients) are: a change in sense of taste, headache, a tingling sensation, feeling faint, indigestion, feeling hot, and excessive sweating.

**Please see additional Important Safety Information on back.
Please see accompanying or click for [Full Prescribing Information](#).**

Can QUZYTIR be used in people of all ages?

Yes, QUZYTIR works in children and adults, including:



Proven safe in children

6 months to 17 years

Children with kidney or liver problems should not take QUZYTIR.



Proven safe in adults, including those

65 years and older

In clinical trials with QUZYTIR, 18 patients were 65 years and older, and 6 patients were 75 years and older. There were no major differences in safety between older and younger patients, but more side effects might be seen in older adults.



Actor portrayals.

SELECT SAFETY INFORMATION

- The most common side effects observed in studies of an oral form of the active ingredient in QUZYTIR taken by adults for a prolonged amount of time were sleepiness, tiredness, dry mouth, sore throat, and dizziness. In similar studies with children, the most common side effects were headache, sore throat, abdominal pain, sleepiness, diarrhea, nosebleed, bronchial spasm, nausea, and vomiting.

Please see additional Important Safety Information on back.

Please see accompanying or click for [Full Prescribing Information](#).

What are the most common side effects?

In the clinical trial, the following were the most common side effects with QUZYTTR, which **occurred in less than 1% of patients:**

- a change in sense of taste
- a tingling sensation
- indigestion
- headache
- feeling faint
- feeling hot
- excessive sweating

QUZYTTR may cause sleepiness. Use caution when driving a car or operating machinery. Avoid alcohol or taking certain other medicines with QUZYTTR, as this may raise your risk of problems with alertness or thinking.

Support is available

QUZYTTR Co-pay Savings Program

Eligible patients may lower their co-pay **to as little as \$0.***



Not actual card.

*For eligible commercially insured patients, the co-pay card is subject to an annual limit. Patients are not eligible if prescriptions are paid for by any state or federally funded program, including, but not limited to, Medicare or Medicaid, Medigap, VA or DOD or TRICARE, or where prohibited by law. Additional terms and conditions apply.

Comprehensive support to help you access treatment

- Helps you understand what financial assistance may be available
- Offers the Patient Assistance Program to eligible patients with established financial need to receive free product. Terms and conditions apply.



For questions or assistance, call **1-866-QUZYTTR (789-9884)**

This is your Q: Ask your doctor if QUZYTIR is right for you

Quzyttir[®]
Cetirizine HCl Injection
10 mg/mL

Scan the QR code to learn more and download
a Doctor Discussion Guide for your next appointment.



What is QUZYTIR?

QUZYTIR is an FDA-approved prescription medicine for the treatment of short-term hives (known as acute urticaria) in adults and children 6 months of age and older.

QUZYTIR is not recommended in children younger than 6 years old with kidney or liver problems.

IMPORTANT SAFETY INFORMATION

- You should not receive QUZYTIR if you are allergic to any of its ingredients or if you are allergic to other medicines that contain levocetirizine or hydroxyzine.
- QUZYTIR may cause sleepiness. Be careful when driving a car or operating potentially dangerous machinery. Avoid alcohol or taking certain other medicines with QUZYTIR, as this may raise your risk of having problems with alertness or thinking. Tell the healthcare provider treating you about all the medicines you are taking, including prescription or over-the-counter medicines, vitamins, or herbal supplements.
- QUZYTIR was not studied in pregnant or nursing women. Tell your healthcare provider if you are pregnant, plan to become pregnant, or are breastfeeding.
- The most common side effects (which occurred in <1% of patients) are: a change in sense of taste, headache, a tingling sensation, feeling faint, indigestion, feeling hot, and excessive sweating.
- The most common side effects observed in studies of an oral form of the active ingredient in QUZYTIR taken by adults for a prolonged amount of time were sleepiness, tiredness, dry mouth, sore throat, and dizziness. In similar studies with children, the most common side effects were headache, sore throat, abdominal pain, sleepiness, diarrhea, nosebleed, bronchial spasm, nausea, and vomiting.
- Call your healthcare provider about any side effects that you may experience.

You are encouraged to report negative side effects of prescription drugs to the FDA. To report suspected adverse reactions, contact the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also contact TerSera Therapeutics at 1-844-334-4035 or medicalinformation@tersera.com.

Please see accompanying or click for [Full Prescribing Information](#).

QUZYTIR, TERSERA, and the TERSERA LOGO are registered trademarks of TerSera Therapeutics LLC or its affiliates.
©2024 TerSera Therapeutics LLC. All rights reserved.
QYR-P-0193 (12/2024)

 **TerSera**[®]
therapeutics