

Ask if QUZYTIR® is right for you

Quzyttir®
Cetirizine HCl Injection
10 mg/mL

When talking with your doctor, use your answers to the following questions to help guide your discussion:

1. What treatments have you received for short-term hives? What has your experience with those treatments been like?

2. Do you have any concerns about your current treatment? If so, please describe.

3. Have you experienced any side effects after receiving treatment for short-term hives? If yes, how would you describe these side effects?

4. What are your treatment goals when you are experiencing short-term hives?

What is QUZYTIR?

QUZYTIR is an FDA-approved prescription medicine for the treatment of short-term hives (known as acute urticaria) in adults and children 6 months of age and older.

QUZYTIR is not recommended in children younger than 6 years old with kidney or liver problems.

SELECT SAFETY INFORMATION

- You should not receive QUZYTIR if you are allergic to any of its ingredients or if you are allergic to other medicines that contain levocetirizine or hydroxyzine.

Please see additional Important Safety Information on the next page.

Please see accompanying [Full Prescribing Information](#).

QUZYTIR has you covered with the next-generation IV antihistamine that lasts 24 hours

Quzyttir[®]
Cetirizine HCl Injection
10 mg/mL

Reasons to ask about QUZYTIR



Treats hives from
allergic reactions



Shown to cause
less drowsiness*



Lasts for
24 hours



Preservative-free
solution for IV use



No major
interactions with
other medications

QUZYTIR is an IV medicine that must be administered by a medical professional.

*Based on a clinical study comparing QUZYTIR to IV diphenhydramine.

SELECT SAFETY INFORMATION

- QUZYTIR may cause sleepiness. Be careful when driving a car or operating potentially dangerous machinery. Avoid alcohol or taking certain other medicines with QUZYTIR, as this may raise your risk of having problems with alertness or thinking. Tell the healthcare provider treating you about all the medicines you are taking, including prescription or over-the-counter medicines, vitamins, or herbal supplements.

Support is available when you're ready



QUZYTIR Reimbursement Resources
for both providers and patients.
To learn more, visit [QUZYTIR.com](https://www.quzyttir.com).



Not actual card

QUZYTIR Co-Pay Program

Eligible patients could lower their co-pay to as little as \$0.†
Learn more at [QUZYTIR.com](https://www.quzyttir.com).

†For eligible, commercially insured patients, the co-pay card is subject to an annual limit. Patients are not eligible if prescriptions are paid by any state or federally funded program, including, but not limited to, Medicare or Medicaid, Medigap, VA, DOD, or TRICARE, or where prohibited by law. Additional terms and conditions apply.

IMPORTANT SAFETY INFORMATION

- You should not receive QUZYTIR if you are allergic to any of its ingredients or if you are allergic to other medicines that contain levocetirizine or hydroxyzine.
- QUZYTIR may cause sleepiness. Be careful when driving a car or operating potentially dangerous machinery. Avoid alcohol or taking certain other medicines with QUZYTIR, as this may raise your risk of having problems with alertness or thinking. Tell the healthcare provider treating you about all the medicines you are taking, including prescription or over-the-counter medicines, vitamins, or herbal supplements.
- QUZYTIR was not studied in pregnant or nursing women. Tell your healthcare provider if you are pregnant, plan to become pregnant, or are breastfeeding.
- The most common side effects (which occurred in <1% of patients) are: a change in sense of taste, headache, a tingling sensation, feeling faint, indigestion, feeling hot, and excessive sweating.
- The most common side effects observed in studies of an oral form of the active ingredient in QUZYTIR taken by adults for a prolonged amount of time were sleepiness, tiredness, dry mouth, sore throat, and dizziness. In similar studies with children, the most common side effects were headache, sore throat, abdominal pain, sleepiness, diarrhea, nosebleed, bronchial spasm, nausea, and vomiting.
- Call your healthcare provider about any side effects that you may experience.

You are encouraged to report negative side effects of prescription drugs to the FDA. To report suspected adverse reactions, contact the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also contact TerSera Therapeutics at 1-844-334-4035 or medicalinformation@tersera.com.

Please see accompanying [Full Prescribing Information](#).