



If your carcinoid syndrome diarrhea relief stalls,
ask your doctor if

**IT'S TIME
FOR XERMELO®**



Indication

XERMELO is a prescription pill, used along with somatostatin analog (SSA) therapy, for carcinoid syndrome diarrhea in adults who are not adequately controlled by SSA therapy.

Select Important Safety Information

- Do not take XERMELO if you have a history of hypersensitivity to telotristat or any of the ingredients in XERMELO.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).

What causes CSD?

Carcinoid syndrome diarrhea (CSD) occurs when a neuroendocrine tumor (NET) cell produces too much serotonin.

Let's take a closer look

To better understand the cause of CSD, take a moment to learn about the science behind it.

1



When cells grow abnormally, they form a tumor.

2



When a tumor forms in the neuroendocrine system, it may produce too many hormones, such as serotonin.

3



Hormones are one way your body's cells communicate with each other to help with bodily functions. For example, serotonin is a hormone that communicates with your gut to help with digestion.

4



When NETs produce too much serotonin, your body's natural serotonin levels can be thrown off balance. The extra serotonin in the gut can lead to CSD.

Select Important Safety Information

- XERMELO may cause constipation which can be serious. You should stop taking XERMELO if severe constipation or severe, persistent, or worsening abdominal pain develops. Talk to your doctor if you have these symptoms.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).

When SSA injections alone are not enough, ask your doctor if it's time for XERMELO

Even though you are being treated with somatostatin analog (SSA) injections, you can still experience symptoms of CSD. **That's why your doctor may add XERMELO**, a prescription pill, to your treatment plan.

Talk to your healthcare provider if you are:



**TAKING
ANTIDIARRHEALS
MORE OFTEN**



**LIMITING ACTIVITIES
AND WORK
OUTSIDE OF HOME**



**PLANNING WHERE TO
FIND BATHROOMS WHEN
YOU LEAVE THE HOUSE**



Actor portrayals.


Select Important Safety Information

- The most common side effects of XERMELO include nausea, headache, increase in liver enzymes, depression, flatulence, decreased appetite, swelling of your hands and feet, fever, abdominal pain, and constipation.

Reduce your serotonin and manage CSD

XERMELO pills + SSA injections work together to reduce serotonin and manage carcinoid syndrome diarrhea (CSD)

SSA blocks release of serotonin outside the cell



XERMELO reduces overproduction of serotonin inside the cell

Unlike antidiarrheal medications, **XERMELO targets a key cause of CSD**, which is too much serotonin in the gut.

How XERMELO was studied

In the TELESTAR study, XERMELO + SSA was compared with SSA injections alone in 90 people (45 per test group) with CSD. Participants were having between 4 to 12 BMs per day and were already on a steady SSA dose for at least 3 months. All participants were followed for 12 weeks and recorded diarrhea frequency in a daily diary.

XERMELO pills + SSA injections provide long-term relief for CSD

TELESTAR, the clinical study that led to the FDA approval of XERMELO, showed meaningful reductions in CSD symptoms.

2X

XERMELO + SSA reduced average daily bowel movement (BM) frequency 2X more than SSA injections alone (-1.4 vs -0.6 BMs/day, respectively).

Select Important Safety Information

- Talk to your doctor about all medications you are taking as some may interact with XERMELO.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).

Give XERMELO 12 weeks to work

In the TELESTAR study, almost 9 out of 10 people with CSD (89%) taking XERMELO pills plus SSA injections had a reduction in diarrhea frequency at 12 weeks. In the same study, only 69% of patients on SSA injections alone reduced their bowel movement frequency.



Most patients see **improvement in 12 weeks** and some patients may **start to see results within 3 weeks**.



Actor portrayal.

Select Important Safety Information

- XERMELO is not recommended if you have moderate or severe liver impairment.

XERMELO reduced serotonin levels

Carcinoid syndrome diarrhea (CSD) occurs when there is too much serotonin in the gut. When serotonin breaks down in your body, it becomes **5-hydroxyindoleacetic acid (5-HIAA)**. **People with carcinoid syndrome diarrhea often have high levels of 5-HIAA.** Your doctor may measure your 5-HIAA to see how much serotonin is in your body.



XERMELO was shown to be **effective** in patients **regardless of their baseline 5-HIAA levels.**

XERMELO was well tolerated in most patients

Let your doctor know about any side effects you experience. The most common side effects of XERMELO include nausea, headache, increase in liver enzymes, depression, flatulence, decreased appetite, swelling of your hands and feet, fever, abdominal pain, and constipation.

In a 12-week clinical study, most patients who started on XERMELO stayed on XERMELO.

Select Important Safety Information

- Do not take XERMELO if you have a history of hypersensitivity to telotristat or any of the ingredients in XERMELO.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).

How to **get started** with XERMELO

You need to take XERMELO 3 times a day, every day, with a meal or snack. That's because food increases your body's ability to absorb XERMELO.



Easy and convenient packaging – first dose and every dose

When it comes to managing your CSD, XERMELO helps make it easy to remember and track every dose.



XERMELO comes in a monthly pack with 4 convenient blister packs (one per week).

For illustration only.



**SYMPTOMS MAY RETURN
IF YOU MISS A DOSE OF XERMELO.**

Select Important Safety Information

- XERMELO may cause constipation which can be serious. You should stop taking XERMELO if severe constipation or severe, persistent, or worsening abdominal pain develops. Talk to your doctor if you have these symptoms.

Get support as you start XERMELO

TerSera NurseSupport is here to support you and serve as a resource for:



Starting XERMELO



Nutrition education



Advocacy connections



CSD education


Once you register, your nurse will **connect with you via phone**. In addition, TerSera NurseSupport participants can receive a nutrition and recipe guide designed to offer CSD-friendly food options.



Enroll in TerSera NurseSupport today!

Scan the **QR code** or visit [TheNetNurse.com](https://www.thenetnurse.com).

The “Find Food You Love” nutrition and recipe guide includes more than 100 pages of easy-to-follow dietary information and recipes designed to **help people with CSD**.



Select Important Safety Information

- The most common side effects of XERMELO include nausea, headache, increase in liver enzymes, depression, flatulence, decreased appetite, swelling of your hands and feet, fever, abdominal pain, and constipation.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).

Financial assistance from TerSera is available



TerSera
SupportSource

- Helps you understand **what financial assistance may be available**
- Offers the Patient Assistance Program to eligible patients with **established financial need** to receive **free product**. Terms and conditions apply



Not actual card.

- Commercially insured patients may **pay as little as \$0 co-pay*** (subject to an annual limit)

*For eligible commercially insured patients, the co-pay card is subject to an annual limit. You are not eligible if prescriptions are paid for by any state or other federally funded programs, including, but not limited to, Medicare or Medicaid, Medigap, VA or DOD or TriCare, or where prohibited by law.



Actor portrayals.

Select Important Safety Information

- Talk to your doctor about all medications you are taking as some may interact with XERMELO.

Indication

XERMELO is a prescription pill, used along with somatostatin analog (SSA) therapy, for carcinoid syndrome diarrhea in adults who are not adequately controlled by SSA therapy.

Important Safety Information

- Do not take XERMELO if you have a history of hypersensitivity to telotristat or any of the ingredients in XERMELO.
- XERMELO may cause constipation which can be serious. You should stop taking XERMELO if severe constipation or severe, persistent, or worsening abdominal pain develops. Talk to your doctor if you have these symptoms.
- The most common side effects of XERMELO include nausea, headache, increase in liver enzymes, depression, flatulence, decreased appetite, swelling of your hands and feet, fever, abdominal pain, and constipation.
- Talk to your doctor about all medications you are taking as some may interact with XERMELO.
- XERMELO is not recommended if you have moderate or severe liver impairment.

You are encouraged to report negative side effects of prescription drugs to the FDA. To report suspected adverse reactions, contact the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also contact TerSera Therapeutics at 1-844-334-4035 or medicalinformation@tersera.com.

Please see accompanying or click for [Full Prescribing Information](#).



Ask your doctor if



For more information, **visit [XERMELO.com](https://www.xermelo.com)**



If your carcinoid syndrome diarrhea (CSD) relief stalls on somatostatin analog (SSA) injections alone, ask your doctor if

**IT'S TIME
FOR XERMELO**

- ✓ Excess serotonin is a key cause of CSD
- ✓ SSA injections may be less effective over time, causing CSD symptoms to return
- ✓ XERMELO pills + SSA injections can help reduce serotonin to help manage CSD
- ✓ XERMELO pills + SSA injections reduced average daily bowel movements 2x more than SSA injections alone
- ✓ When starting XERMELO, it may take 12 weeks to see full improvement. Others may see improvement in as little as 3 weeks
- ✓ TerSera offers you resources with its NurseSupport and SupportSource programs

Indication

XERMELO is a prescription pill, used along with somatostatin analog (SSA) therapy, for carcinoid syndrome diarrhea in adults who are not adequately controlled by SSA therapy.

Select Important Safety Information

- Do not take XERMELO if you have a history of hypersensitivity to telotristat or any of the ingredients in XERMELO.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).