



If your carcinoid syndrome diarrhea relief stalls,

**IT'S TIME
FOR XERMELO**



Indication

XERMELO is a prescription pill, used along with somatostatin analog (SSA) therapy, for Carcinoid Syndrome diarrhea in adults who are not adequately controlled by SSA therapy.

Select Important Safety Information

- Do not take XERMELO if you have a history of hypersensitivity to telotristat or any of the ingredients in XERMELO.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).


What causes CSD?


Carcinoid syndrome diarrhea (CSD) occurs when a neuroendocrine tumor (NET) cell produces too much serotonin.


Let's take a closer look

To better understand the cause of CSD, take a moment to learn about the science behind it.

1  **When cells grow abnormally,** they form a tumor.

2  **When a tumor forms** in the neuroendocrine system, it may produce too many hormones, such as serotonin.

3  **Hormones are one way** your body's cells communicate with each other to help with bodily functions. For example, serotonin is a hormone that communicates with your gut to help with digestion.

4  **When NETs produce too much serotonin,** your body's natural serotonin levels can be thrown off balance. The extra serotonin in the gut can lead to CSD.

When SSA injections alone are not enough, it's time for XERMELO

Even though you are being treated with somatostatin analog (SSA) injections, you can still experience symptoms of CSD. **That's why you and your doctor may add XERMELO,** a prescription pill, to your treatment plan.

Talk to your healthcare provider if you are:



**TAKING
ANTI-DIARRHEALS
MORE OFTEN**



**LIMITING ACTIVITIES
AND WORK
OUTSIDE OF HOME**



**PLANNING WHERE TO
FIND BATHROOMS WHEN
YOU LEAVE THE HOUSE**



Actor portrayals.

Select Important Safety Information

- XERMELO may cause constipation which can be serious. You should stop taking XERMELO if severe constipation or severe, persistent, or worsening abdominal pain develops. Talk to your doctor if you have these symptoms.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).

Select Important Safety Information

- The most common side effects of XERMELO include nausea, headache, increase in liver enzymes, depression, flatulence, decreased appetite, swelling of your hands and feet, fever, abdominal pain, and constipation.

Reduce your serotonin and manage CSD

XERMELO pills + SSA injections work together to reduce serotonin and manage CSD



Unlike antidiarrheal medications, **XERMELO targets the cause of CSD**, which is too much serotonin in the gut.

XERMELO pills + SSA injections provide long-term relief for CSD

TELESTAR, a major clinical study that led to the FDA approval of XERMELO, showed meaningful reductions in CSD symptoms.



XERMELO + SSA reduced daily bowel movement (BM) frequency **2X** more than SSA injections alone (-1.4 vs -0.6 BMs/day, respectively).

How XERMELO was studied

In the TELESTAR study, XERMELO + SSA was compared with SSA injections alone in people with carcinoid syndrome. All people were followed for 12 weeks and recorded diarrhea frequency in a daily diary.

Select Important Safety Information

- Talk to your doctor about all medications you are taking as some may interact with XERMELO.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).

Give XERMELO 12 weeks to work

In a 12-week study, almost 9 out of 10 people with CSD (89%) taking XERMELO pills plus SSA injections had a reduction in diarrhea frequency. In the same study, only 69% of patients on SSA injections alone reduced their bowel movement frequency.



Most patients see **significant improvement in 12 weeks** and some patients may **start to see results within 3 weeks**.



Actor portrayal.

Select Important Safety Information

- XERMELO is not recommended if you have moderate or severe liver impairment.

XERMELO reduced serotonin levels

CSD occurs when there is too much serotonin in the gut. When serotonin breaks down in your body, it becomes **5-hydroxyindoleacetic acid (5-HIAA)**. **People with carcinoid syndrome often have high levels of 5-HIAA.** Your doctor may measure the 5-HIAA to see how much serotonin is in your body.



XERMELO was shown to be **effective** in patients **regardless of their baseline 5-HIAA levels.**

XERMELO was safe and well tolerated in most patients

Let your doctor know about any side effects you experience. Most patients who started on XERMELO stayed on XERMELO.

Select Important Safety Information

- Do not take XERMELO if you have a history of hypersensitivity to telotristat or any of the ingredients in XERMELO.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).

How to get started with XERMELO

You need to take XERMELO 3 times a day, every day, with a meal or snack. That's because food increases your body's ability to absorb XERMELO.



Easy and convenient – first dose and every dose

When it comes to managing your CSD, XERMELO helps make it easy to remember and track every dose.



XERMELO comes in a monthly pack with 4 convenient blister packs (one per week).

For illustration only.



**SYMPTOMS MAY RETURN
 IF YOU MISS A DOSE OF XERMELO.**

Select Important Safety Information

- XERMELO may cause constipation which can be serious. You should stop taking XERMELO if severe constipation or severe, persistent, or worsening abdominal pain develops. Talk to your doctor if you have these symptoms.

Get support as you start XERMELO

TerSera NurseSupport is here to support you and provide recommendations to help you live your best life with CSD. Your nurse can serve as a resource for:



Starting XERMELO



Nutrition education



Advocacy connections



CSD education


Once you register, your nurse will **connect with you via phone**. In addition, TerSera NurseSupport participants can receive a nutrition and recipe guide designed to offer carcinoid syndrome-friendly food options.



Enroll in TerSera NurseSupport today!

Scan the **QR code** or visit TheNetNurse.com.

The “Find Food You Love” nutrition and recipe guide includes more than 100 pages of easy-to-follow dietary information and recipes designed to **help people with CSD**.




Select Important Safety Information

- The most common side effects of XERMELO include nausea, headache, increase in liver enzymes, depression, flatulence, decreased appetite, swelling of your hands and feet, fever, abdominal pain, and constipation.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).

Financial assistance from TerSera is available



Not actual card.

- Commercially insured patients may **pay as little as \$0* co-pay** up to \$10,000/year per patient



TerSera
SupportSource

- Helps you **understand what financial assistance is available** through foundations and other third-party organizations
- Offers the Patient Assistance Program to those with **established financial need** to receive **free product**

*For eligible commercially insured patients, card carries a maximum annual benefit of \$10,000 per calendar year. You are not eligible if prescriptions are paid by any state or other federally funded programs, including, but not limited to Medicare or Medicaid, Medigap, VA or DOD or TriCare, or where prohibited by law.



Actor portrayals.

Select Important Safety Information

- Talk to your doctor about all medications you are taking as some may interact with XERMELO.

Indication

XERMELO is a prescription pill, used along with somatostatin analog (SSA) therapy, for Carcinoid Syndrome diarrhea in adults who are not adequately controlled by SSA therapy.

Important Safety Information

- Do not take XERMELO if you have a history of hypersensitivity to telotristat or any of the ingredients in XERMELO.
- XERMELO may cause constipation which can be serious. You should stop taking XERMELO if severe constipation or severe, persistent, or worsening abdominal pain develops. Talk to your doctor if you have these symptoms.
- The most common side effects of XERMELO include nausea, headache, increase in liver enzymes, depression, flatulence, decreased appetite, swelling of your hands and feet, fever, abdominal pain, and constipation.
- Talk to your doctor about all medications you are taking as some may interact with XERMELO.
- XERMELO is not recommended if you have moderate or severe liver impairment.

You are encouraged to report negative side effects of prescription drugs to the FDA. To report suspected adverse reactions, contact the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also contact TerSera Therapeutics at 1-844-334-4035 or medicalinformation@tersera.com.

Please see accompanying or click for [Full Prescribing Information](#).

PI



Talk to your doctor when



For more information, visit XERMELO.com



If your carcinoid syndrome diarrhea relief stalls,

**IT'S TIME
FOR XERMELO**

- ✓ Excess serotonin is the root cause of carcinoid syndrome diarrhea, or CSD
- ✓ SSA injections may be less effective over time, causing CSD symptoms to return
- ✓ XERMELO pills + SSA injections work together to reduce serotonin to help manage CSD
- ✓ XERMELO pills + SSA injections reduced daily bowel movements 2x more than SSA injections alone
- ✓ When starting XERMELO, it may take 12 weeks to see full improvement. Others may see improvement in as little as 3 weeks
- ✓ TerSera offers you resources with its NurseSupport and SupportSource programs

Indication

XERMELO is a prescription pill, used along with somatostatin analog (SSA) therapy, for Carcinoid Syndrome diarrhea in adults who are not adequately controlled by SSA therapy.

Select Important Safety Information

- Do not take XERMELO if you have a history of hypersensitivity to telotristat or any of the ingredients in XERMELO.

Please see Full Important Safety Information on page 10.

Please see accompanying or click for [Full Prescribing Information](#).